

Testing population synthesis models with globular cluster colors

Pauline Barmby & John P. Huchra

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

ABSTRACT

We have measured an extensive set of *UBVRIJHK* colors for M31 globular clusters (Barmby et al. 2000). We compare the predicted simple stellar population colors of three population synthesis models to the intrinsic colors of Galactic and M31 globular clusters. The best-fitting models fit the cluster colors very well – the weighted mean color offsets are all < 0.05 mag. The most significant offsets between model and data are in the *U* and *B* passbands; these are not unexpected and are likely due to problems with the spectral libraries used by the models. The metal-rich clusters ($[\text{Fe}/\text{H}] \gtrsim -0.8$) are best fit by young (8 Gyr) models, while the metal-poor clusters are best fit by older (12–16 Gyr) models. If this range of globular cluster ages is correct, it implies that conditions for cluster formation must have existed for a substantial fraction of the galaxies’ lifetimes.

Subject headings: galaxies: individual (M31) – galaxies: star clusters – globular clusters: general – galaxies: stellar content

1. Introduction

Predicting the integrated spectral energy distributions of stellar populations is important in the solution of many problems in astronomy, from determining the ages of globular clusters to modeling counts of faint galaxies at high redshift. Beginning with the early work of Tinsley (1968), successive generations of modelers have combined the best available data on stellar structure and evolution to predict the appearance of the combined light of generations of stars. Although the subject of population synthesis has a long history, it is an active area of research: synthesis techniques and many of the input data (isochrones, opacities, spectral libraries) continue to be improved.

There is good evidence that globular clusters (GCs) are internally homogeneous in age and metallicity (Heald et al. 1999; Stetson 1993). GCs are the best observational analogs of modelers’ ‘simple stellar populations’, i.e. populations of stars formed over a short time out of gas with homogeneous chemical composition. Broadband colors are among the simplest predictions of population synthesis models, so comparing the models’ predicted colors to cluster colors is the natu-

ral zeroth-order test of compatibility between the models and reality (Huchra 1996). In this paper we compare the broad-band *UBVRIJHK* colors predicted by three modern population synthesis models with the colors of Galactic and M31 GCs. Our observational database is the first one to include extensive coverage of the *JHK* bandpasses, and the first with spectroscopic metallicities for all clusters. We use the cluster metallicities to bin the clusters for comparison to the appropriate models. In this way we determine the cluster-to-model offset separately for each color and avoid the ambiguity in comparing model and cluster colors in two-color diagrams.

2. Input data and comparison procedure

For M31 clusters, the observational data are from the Barmby et al. (2000) catalog. For the Galactic clusters we obtained optical colors, metallicity, and reddening from the June 1999 version of the Harris (1996) catalog, and IR colors from Frogel et al. (1980), as reported in Brodie & Huchra (1990) (but with the reddening correction applied by Brodie & Huchra (1990) removed and the reddening values in Harris (1996)

used instead). We dereddened the clusters' colors using the values of E_{B-V} given in the catalogs and the Cardelli et al. (1989) extinction curve for $R_V = 3.1$. For M31, we excluded clusters where the error in the spectroscopic metallicity was $\sigma_{[\text{Fe}/\text{H}]} > 0.5$, and clusters suspected of being young on the basis of strong Balmer absorption or blue $B-V$ colors (see Barmby et al. 2000). For both galaxies, we excluded clusters with $E_{B-V} > 0.5$; there are 103 M31 and 85 Galactic clusters in the final sample. Photometric data is not available in all bandpasses for all clusters: only about two-thirds have measured R and I , and less than half have H .

We compare the cluster colors to those for simple stellar populations of ages 8, 12, and 16 Gyr from three sets of models: those of Worthey¹, Bruzual and Charlot (hereafter BC) (both the Worthey and BC models are reported in Leitherer et al. 1996), and Kurth et al. (1999) (hereafter KFF). Although model colors are tabulated in smaller age increments (typically 1 Gyr), initially it is more reasonable to use the models as a rough guide to relative ages rather than attempting to derive precise cluster ages from them. The Worthey models are computed at $[\text{Fe}/\text{H}]$ values of $-2.0, -1.5, -1.0, -0.5$, and -0.25 dex, and the BC and KFF models are computed at $[\text{Fe}/\text{H}]$ values of -2.33 (KFF models only), -1.63 , -0.63 , and -0.32 dex. We compared clusters to both the Salpeter IMF (Worthey's 'vanilla' models) and Scalo (1986) (Miller & Scalo 1979, in the Worthey models) IMF version of the models. Worthey (1994) finds that some of his model colors have defects (e.g. $B-V$ is too red by $0.04-0.06^m$ due to problems in the theoretical stellar atmospheres and the color-temperature calibration), but the sizes of these defects are not well-determined so we do not correct for them. Figure 1 shows data and models in two frequently-used two-color diagrams.

Since the models are computed at discrete values of $[\text{Fe}/\text{H}]$, we use the spectroscopic metallicities of the clusters to compare only clusters with comparable metallicities (± 0.25 dex) to each model. The Galactic cluster metallicities given in

Harris (1996) are on the Zinn & West (1984) (ZW) metallicity scale, and the M31 cluster metallicities are also tied to this scale through the calibration of Brodie & Huchra (1990). Recent work (Carretta & Gratton 1997; Rutledge et al. 1997) suggests that the ZW scale may be non-linear at both high and low metallicities. We retain the ZW scale in this paper because we found that using the Carretta & Gratton (1997) scale to assign clusters to model comparison bins made little difference in our results. We caution, however, that the effect of changing the metallicity scale is unknown for the $[\text{Fe}/\text{H}] = -0.25$ model bin. The transformation from the ZW to CG scales is only defined for $[\text{Fe}/\text{H}]_{\text{ZW}} < -0.5$, the lower limit of this metallicity bin.

We calculated the mean offsets between model and cluster colors (referenced to V) for each metallicity bin; Figures 2–3 show some representative comparisons. We plot $\Delta(X-V)$ for all bandpasses X to make clear the differences in spectral energy distributions between models and data; we remind the reader that the offsets for bandpasses redward of V thus have the opposite sign from the usual colors. One general characteristic of the models visible in the Figures is that younger-aged models predict bluer colors. The exception is the KFF Scalo model for $[\text{Fe}/\text{H}] = -1.63$, which predicts only very small color differences ($\lesssim 0.01^m$) between ages of 12 and 16 Gyr. The effect of the IMF on the colors appears to depend on both metallicity and age. For the Worthey $[\text{Fe}/\text{H}] = -1.50$ models, Miller-Scalo IMF colors are redder than Salpeter model colors at all ages, but for the $[\text{Fe}/\text{H}] = -0.50$ models, the Miller-Scalo IMF colors are bluer for 8 and 12 Gyr and almost identical for 16 Gyr. BC predict almost no color difference between the Salpeter and Scalo IMF models of the same age and metallicity.

A striking feature in Figures 2 and 3 is the range of discrepancies between models and data. For example, the largest difference between the Worthey model with parameters (Salpeter IMF, $[\text{Fe}/\text{H}] = -1.50$, age 16 Gyr) and the mean colors of clusters with $-1.75 \leq [\text{Fe}/\text{H}] \leq -1.25$ is 0.04^m in $U-V$. The same models with $[\text{Fe}/\text{H}] = -0.50$ are well offset from the data at all colors except $U-V$; the largest offset is 0.23^m in $V-K$. To determine the best-fitting models, we quantify the overall goodness-of-fit for each model/cluster

¹The version we used updates the Worthey (1994) models by including a more realistic treatment of the horizontal branch for $[\text{Fe}/\text{H}] < -1.0$.

metallicity bin pair as:

$$F = \frac{\sum_k |\Delta(X - V)_k| / \sigma_k^2}{\sum_k 1 / \sigma_k^2} \quad (1)$$

The color differences $\Delta(X - V)_k$ are weighted by $1/\sigma_k^2$, where σ_k are the standard errors in the mean colors of objects in the bin. Table 1 gives the Δ and F values for the best fitting models in each metallicity bin.

3. Discussion

Table 1 shows that the best-fitting models fit the data quite well, with typical color offsets of $0.02 - 0.03^m$. The two bandpasses with the most significant offsets are U and B : the models are too blue in $U - V$ and too red in $B - V$. Neither offset shows a clear trend with metallicity. The offsets are likely not due to systematics in the photometric system or in the extinction curve. While problems with the photometric systems might be expected in the R and I bands (due to conversion between the Johnson and Cousins RI systems), both data and models use the well-defined Johnson UBV system. Problems in the reddening curve also seem unlikely for the same reasons. We suspect that the offsets are more likely due to systematic errors in the models. The $B - V$ offset in particular is likely due to the flux libraries used. Both Worthey (1994) and Lejeune (1997) found their model $B - V$ colors to be $0.04 - 0.06^m$ too red compared to empirical solar-metallicity spectra, even after correcting to the empirical color-temperature scale. This suggests a possible problem with the stellar atmosphere models of Kurucz (1995), upon which both libraries are based.

The cause of the offset in $U - V$ is not as clear. This offset is actually worse than it appears: since we compute the $U - V$ colors for the BC and KFF models as $(U - B) + (B - V)$, the red $B - V$ colors *compensate* for some of the $U - B$ defect, which is actually larger than the defect in $U - V$. Worthey (1994) – whose models give $U - V$ directly – finds that his model $U - V$ is too blue compared to solar neighborhood stars and elliptical galaxies. Worthey cites problems with the U fluxes from the stellar libraries as a possible cause: modeling the many blended atomic and molecular lines blueward of B is difficult, and many of the necessary opacities are not well determined. This

cannot be the only cause of model problems in U , since the BC and KFF models, which use the same stellar library, predict different $U - V$ colors. The treatment of the horizontal branch in the models is another possible source of problems in the $U - V$ colors because the HB emits most of the blue light. However, systematic problems with the model HB color (which depends on metallicity), would presumably produce a $U - V$ offset dependent on metallicity, which we do not observe. Observational error is another possible contributor to the $U - V$ offset, as many of the $U - B$ colors of the M31 clusters are poorly determined (see Table 3 of Barmby et al. 2000). Understanding the rest-frame U flux of stellar populations becomes increasingly important when studying high-redshift galaxies and global star formation history, and further investigation of the models in this bandpass is clearly warranted.

A secondary result in Table 1 is that age determines which model best fits the data. Higher-metallicity cluster colors are best fit by 8 Gyr models, regardless of IMF. Lower-metallicity cluster colors ($[\text{Fe}/\text{H}]_{\text{bin}} \leq -1.00$) are best fit by 12 or 16 Gyr models. The best-fit age depends on the IMF for several of the models, but not in any systematic fashion. This result is consistent with the determinations of relative ages for Galactic clusters by Rosenberg et al. (1999). These authors determined relative ages of 35 Galactic globular clusters from a homogeneous set of V , $V - I$ color-magnitude diagrams. They compared theoretical isochrones with the observational CMDs to determine ages using two independent methods. They found that the clusters with $[\text{Fe}/\text{H}]_{\text{CG}} > -0.9$ were $\sim 17\%$ younger than clusters with $[\text{Fe}/\text{H}]_{\text{CG}} < -1.2$, with the intermediate-metallicity clusters showing a $\sim 25\%$ age dispersion. These results are model-dependent, as are ours, but the results' similarity implies that either there is a real difference between metal-rich and metal-poor clusters or there is a systematic problem in the models in one of the metallicity regimes.

What possible systematic errors in our input data or comparison procedure could produce the result that the metal-rich clusters are younger? We redid the comparison procedure considering the clusters of each galaxy separately, and still found younger ages for the most metal-rich clus-

ters. Although M31 has a greater proportion of the metal-rich clusters, younger ages are found for both M31 and Galactic metal-rich clusters. Cohen & Matthews (1994) suggest that the spectroscopic metallicities of the most metal-rich M31 clusters measured by Huchra et al. (1991) are systematically too high. If this is true, the clusters would appear too blue compared to old, higher-metallicity models and the best-fit model would be younger. We compared the metal-rich M31 clusters to the Worthey $[\text{Fe}/\text{H}] = -1.0$ models, and the best-fitting model had age 16 Gyr. However, the goodness-of-fit was better for the young, metal-rich models than for the older, more metal-poor model, so we conclude that younger ages are still favored for these clusters. Overestimating the reddening of the metal-rich clusters would make the derived intrinsic colors too blue and yield younger ages. This seems unlikely, given that the color-metallicity relations for Galactic and M31 clusters match well throughout their metallicity range (see Barmby et al. 2000), and the methods of reddening determination for M31 and Galactic clusters are different.

If the detection of younger ages for metal-rich globular clusters is real, it has implications for galaxy formation. A range of GC ages implies that GC formation took place over an extended period of time. Conditions for GC formation were not particular to the early universe, an assertion supported by observations of ‘proto-globular’ clusters in present-day merging galaxies (e.g. Zepf et al. 1999). More precise knowledge of the distribution of cluster ages in each galaxy would be extremely useful in understanding cluster system formation. If the age distribution is continuous, the relation between age and metallicity might hold clues as to what factors controlled the cluster formation rate. If the age distribution is bimodal – with most clusters old and coeval and the remainder younger and coeval – then some event must have triggered the second episode of GC formation. Perhaps the younger clusters were stripped from or accreted along with satellite galaxies of M31 and the Galaxy.

4. Conclusions

Comparison of three sets of population synthesis models with integrated colors of M31 and

Galactic globular clusters shows that the models reproduce the redder average cluster colors to within the observational uncertainties. The poorer agreement in $U-V$ and $B-V$ is likely due to systematic errors in the models. Younger-age models are required to best match the colors of the metal-rich clusters, consistent with the findings of Rosenberg et al. (1999) that the most metal-rich Galactic clusters are younger than the bulk of the globular cluster population. A range of ages for globular clusters implies that conditions for cluster formation were not restricted to the early universe. The cluster age distribution has important implications for galaxy and globular cluster system formation, and attempts to determine it more precisely are needed.

REFERENCES

- Barmby, P., Huchra, J. P., Brodie, J. P., Forbes, D. A., Schroder, L. L., & Grillmair, C. J. 2000, *AJ*, in press (astro-ph/9911152)
- Brodie, J. P. & Huchra, J. P. 1990, *ApJ*, 362, 503
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Carretta, E. & Gratton, R. G. 1997, *A&AS*, 121, 95
- Cohen, J. G. & Matthews, K. 1994, *AJ*, 108, 128
- Frogel, J. A., Persson, S. E., & Cohen, J. G. 1980, *ApJ*, 240, 785
- Harris, W. H. 1996, *AJ*, 112, 1487
- Heald, A. J., Griffiths, W. K., Penny, A. J., & Morris, P. W. 1999, *MNRAS*, 307, 789
- Huchra, J. P. 1996, in *From Stars to Galaxies: the Impact of Stellar Physics on Galaxy Evolution*, ed. C. Leitherer, U. F. von Alvensleben, & J. Huchra (San Francisco: ASP), 597
- Huchra, J. P., Brodie, J. P., & Kent, S. M. 1991, *ApJ*, 370, 495
- Kurth, O. M., Fritze-von Alvensleben, U., & Fricke, K. J. 1999, *A&AS*, 138, 19 (KFF)
- Kurucz, R. 1995, private communication

Lejeune, T. 1997, PhD thesis, Univ. Louis Pasteur, Strasbourg & Astron. Institute of Basel University, Basel

Leitherer, C. et al. 1996, *PASP*, 108, 996

Miller, G. E. & Scalo, J. M. 1979, *ApJS*, 41, 513

Rosenberg, A., Saviane, I., Piotto, G., & Aparicio, A. 1999, *AJ*, 118, 2306

Rutledge, G. A., Hesser, J. E., & Stetson, P. B. 1997, *PASP*, 109, 907

Scalo, J. M. 1986, *Fund. Cosmic Phys.*, 11, 1

Stetson, P. B. 1993, in *The Globular Cluster-Galaxy Connection*, ed. G. H. Smith & J. P. Brodie (San Francisco: ASP), 14

Tinsley, B. 1968, *ApJ*, 151, 547

Worthey, G. 1994, *ApJS*, 95, 107

Zepf, S. E., Ashman, K. M., English, J., Freeman, K. C., & Sharples, R. M. 1999, *AJ*, 118, 752

Zinn, R. & West, M. J. 1984, *ApJS*, 55, 45




Fig. 1.— $(B - V)_0$ vs. $(U - V)_0$ and $(V - K)_0$ for M31 globular clusters (triangles) and Galactic GCs (squares). Lines are population synthesis models of ages 8 Gyr (bluer colors) and 16 Gyr (redder colors): Worthey (solid), BC (dashed), KFF (dotted).




Fig. 2.— Color offsets $\Delta(X - V)$ (data–models) for Salpeter IMF. ($\Delta V = 0$ is plotted to emphasize that the models are normalized to the data at V .) Solid lines: 16 Gyr models, dotted lines: 12 Gyr models, dashed lines: 8 Gyr models. The -1.5 and -0.5 bins in $[\text{Fe}/\text{H}]$ are Worthey models; the -1.63 and -0.63 $[\text{Fe}/\text{H}]$ bins are BC models. Error bars (plotted only on the 8 Gyr models for clarity) are the standard errors of the mean cluster colors and do not include observational uncertainties.




Fig. 3.— Color offsets $\Delta(X - V)$ or $\Delta(V - X)$ (data-models) for Scalo or Miller-Scalo IMF. Symbols same as Figure 2. The -1.5 and -0.5 bins in $[\text{Fe}/\text{H}]$ are Worthey models; the -1.63 and -0.63 $[\text{Fe}/\text{H}]$ bins are KFF models (which have no $\Delta(J - V), \Delta(H - V)$ since they do not predict these colors).

TABLE 1
COLOR OFFSETS FOR BEST-FITTING MODELS

[Fe/H]	model	F	N_{gc}	$\Delta(U - V)$	$\Delta(B - V)$	$\Delta(V - R)$	$\Delta(V - I)$	$\Delta(V - J)$	$\Delta(V - H)$	$\Delta(V - K)$
-0.25	W8sc	0.023	15	23.4 ± 2.4	-0.4 ± 0.9	-3.0 ± 1.1	-0.3 ± 1.0	3.2 ± 2.6	-1.4 ± 6.0	-6.0 ± 4.9
-0.32	K8sc	0.022	17	11.5 ± 2.2	-2.5 ± 0.8	-2.1 ± 1.0	0.2 ± 0.9	...	...	-3.9 ± 4.6
-0.50	W8sc	0.013	30	22.8 ± 3.1	-0.3 ± 0.8	-1.0 ± 0.7	1.2 ± 1.3	4.6 ± 4.6	2.8 ± 5.2	-3.3 ± 7.2
-0.63	K8sp	0.012	35	12.2 ± 3.2	-1.0 ± 0.9	-0.6 ± 0.6	2.4 ± 1.2	...	...	2.1 ± 5.9
-1.00	W12sc	0.014	48	13.7 ± 2.2	-2.3 ± 1.0	0.0 ± 0.5	-1.1 ± 1.2	3.2 ± 1.8	6.9 ± 2.5	4.3 ± 2.1
-1.50	W16sp	0.016	75	4.1 ± 1.4	-3.0 ± 0.6	-0.3 ± 0.5	-1.0 ± 0.8	2.0 ± 1.7	-0.7 ± 3.5	2.0 ± 2.3
-1.63	K16sp	0.019	77	5.1 ± 1.2	-0.7 ± 0.5	1.7 ± 0.4	2.8 ± 0.7	...	...	10.8 ± 2.1
-2.00	W16sp	0.027	37	8.2 ± 1.2	-2.6 ± 0.6	-0.3 ± 0.8	-1.6 ± 1.2	-6.7 ± 2.6	-5.2 ± 3.7	-5.8 ± 3.5
-2.23	K16sp	0.027	12	4.1 ± 2.3	1.8 ± 1.3	2.3 ± 2.1	3.4 ± 2.0	...	...	-7.1 ± 6.2

NOTE.—Units are hundredths of a magnitude. The capital letter in the model column indicates the best-fitting model (Worthey, BC or KFF), the number is the model age in Gyr, and ‘sp’ or ‘sc’ indicate Salpeter or Scalo IMF, respectively.